

Articles of Interest

Literature Review

Reviewers

Heather L. Orsted, RN, BN,
ET, MSc

Leah Shapera, RN, MSN,
GNC(c)

The Paths from Research to Improved Health Outcomes

Authors: Glasziou P, Haynes B

Publication: *Evidence-Based Medicine*. 2005;10:4-7.

Reviewer: Heather L. Orsted, RN, BN, ET, MSc

What happens when McMaster meets Oxford? This is an interesting discussion paper on how evidenced-based medicine (EBM) "should not just be concerned with clinical content but also about the processes of changing care and systems of care." Glasziou and Haynes simply ask clinicians to look at two aspects of EBM: you need to get the evidence *straight* and you need to get the evidence *used*.

One key point in this paper is that the discussion does not end with clinicians getting the information; it also involves getting patients to adhere to practice based on the evidence. This paper also includes a diagram depicting the research-to-practice pipeline, which is appealing for visual learners.

This is a very useful paper for the clinician struggling to support a

shift toward best practice.

Further to this article, Glasziou and Haynes are on the team that has recently published *Evidence-Based Medicine: How to Practice and Teach EBM*, Third Edition.*

*Straus S, Scott Richardson W, Glasziou P, Haynes RB. *Evidence-Based Medicine*, Third Edition. Edinburgh: Elsevier Churchill Livingstone. 2005.

Analgesic Effects of Topical Methadone: A Report of Four Cases

Authors: Gallagher RE, Arndt DR, Hunt KL

Publication: *Clinical Journal of Pain*. 2005;21(2):190-192.

Reviewer: Leah Shapera, RN, MSN, GNC(c)

The objective of this case series was to find both an opioid and a delivery system that would provide lasting pain relief between dressing changes for patients with open wounds. While studies have shown that topical morphine (often mixed with a gel) can be applied to open wounds to achieve a degree of pain relief, there is significant variation reported in the duration of pain relief achieved. This means that relief will not consistently last between daily dressing changes, which makes it generally unsuitable for the palliative care population.

In this study, four cases are presented. In each case, methadone powder (100 mg) mixed in

absorbent protective powder (10 g) is sprinkled on the open wound once daily at the time of each dressing change.

The authors found that the best results were achieved when using the topical preparation on more exudative wounds with exposed tissue. Drier wounds with eschar showed less positive response, as the powder mixture tended to adhere to the wound beds. There were no reported or observed adverse effects from the topical methadone, nor did the mixture interfere with wound healing. As the authors point out, it has been noted that opioids may actually reduce wound inflammation, which would help with both pain and tissue repair.

In the last case presented, the patient did not experience any analgesic effect from the methadone mixture (after multiple attempts). This same patient had no significant improvement with oral methadone either. In this case the authors presumed that the patient did not have opioid receptors sensitive to

methadone, and point out how this illustrates that "peripheral opioid receptors of an individual are from the same population as that individual's central receptors."

In summary, this short case series demonstrates that topical methadone is absorbed and can be effective in controlling pain in exuding wounds with exposed tissue. The degree of topical absorption is variable and likely somewhat dependent upon the site of the wound, the amount of eschar, and the degree of local peri-wound circulation.

It would be important for this study to be replicated with a larger sample size. Additionally, studying the effects of the topical methadone mixture used could be extended to populations other than palliative care. Nonetheless, the authors are to be commended for their innovative efforts in searching for new and more effective methods of pain management, as these are desperately needed to improve the comfort and quality of life for patients with wounds. ☺

**You've Got Questions,
We've Got Answers**

**The CAWC Discussion Forum at
www.cawc.net is the only place
in Canada accessible to every
wound-care clinician in the country.**

**It's a great way to tap into a
national network of wound-care professionals. Visit it today!**



Take a closer look.

The 3M™ Tegaderm™ brand is expanding.

The Tegaderm™ brand you've trusted for over 20 years is expanding its family of dressings to include:

- Advanced Wound Care
- Surgical Site
- IV Site Secural & Protection

New packaging and naming will simplify product selection, and new products are designed to be simple, dependable, and trusted. And be sure to look for Tegaderm Ag Mesh – our newest member of the *Tegaderm Family of Products*.

Visit the new Tegaderm™ brand website at www.3M.com/Tegaderm or call 1-800-563-2921.

3M™ Tegaderm™
Dressings

SIMPLE. DEPENDABLE. TRUSTED.

Product shown is 3M™ Tegaderm™ Transparent Film Dressing #9519HP.
© 3M 2006. All Rights Reserved. 0607-MS-26275

3M Health Care

Palliative Dilemmas *continued from page 13*

6. Beckett R, Coombs TJ, Frost MR, McLeish J, Thompson K. Charcoal cloth and malodorous wounds. *Lancet*. 1980;2(8194):594-595.
7. Price E. Wound care: The stigma of smell. *Nursing Times*. 1996;88(20):71-72.
8. Grocott P. Assessment of fungating malignant wounds. *Journal of Wound Care*. 1995;4(7):333-336.
9. Boardman M, Mellor K, Neville B. Treating a patient with a heavily exudating malodorous fungating ulcer. *Journal of Wound Care*. 1993;8(5):216-218.
10. Langemo DK. Quality of life and pressure ulcers: What is the impact? *WOUNDS*. 2005;17(1):3-7.
11. Fleck CA. Ethical wound management for the palliative patient. *Extended Care Product News*. 2005;100(4):38-46.
12. Moody M. Metrotop: A topical antimicrobial agent for malodorous wounds. *Br J Nurs*. 1998;7:286-289.
13. Hampson JP. The use of metronidazole in the treatment of malodorous wounds. *Journal of Wound Care*. 1996;5(9):421-425.
14. Van Toller S. Invisible wounds: The effects of skin ulcer malodors. *Journal of Wound Care*. 1994;3(2):103-106.
15. Gold GH, Pugh EN. The nose leads the eye. *Nature*. 1997;385:677-679.
16. Dalton P, Wysocki CJ, Brady MJ, Lawley HJ. The influence of cognitive bias on the perceived odor, irritation and health symptoms from chemical exposure. *Int Arch Occup Environ Health*. 1997;69:407-417.
17. Sapico FL, Ginunas VJ, Thornhill-Joyes M, Canawati HN, Capen DA, Klein NE. Quantitative microbiology of pressure sores in different stages of healing. *Diagn Microbiol Infect Dis*. 1986;5:31-38.
18. Lindholm C. Pressure ulcers and infection: Understanding clinical features. *Ostomy/Wound Management*, May 2003;49(5A):4-7.
19. Kingsley A. A proactive approach to wound infection. *Nursing Standard*. 2001;15(30):50-58.
20. Percival SL, Percival PG, Bowler PG, Russell D. Bacterial resistance to silver in wound care. *Journal of Hospital Infection*. 2005;60:1-7.
21. Kalinski C, et al. Effectiveness of a topical formulation containing metronidazole for wound odor and exudate control. *WOUNDS*. 2005;17(4):84-90.
22. Köse AA, Karabagli Y, Kırkçüoğlu M, Çetin C. Alpha Sepiolite: An old clay mineral, a new dressing material. *WOUNDS*. 2005;17(5):114-121.
23. Ovington L. Bacterial toxins and wound healing. *Ostomy Wound Management*. 2003;49(7A):8-12.
24. Lipman RDA. Avery Dennison Medical. Odour absorbing hydrocolloid dressing for direct wound contact [Poster Number 82]. Wound Healing Society (WHS) 15th Annual Meeting, Chicago. May 18-21, 2005.
25. Fleck CA, Frizzell LD. When negative is positive: A review of negative pressure wound therapy. *ECPN*. 2004;92(2):20-25.

Processus de validation d'une traduction française ... *suite de la page 28*

- Scale for predicting pressure sore risk. *Nursing Clinics of North America*. 1987;22(2):417-428.
7. Braden B, Bergstrom N. A conceptual schema for the study of the etiology of pressure sores. *Rehabilitation Nursing*. 1987;12(1):8-12.
8. —. Clinical utility of the Braden Scale for predicting pressure sore risk. *Decubitus*. 1989;2(3):44-46, 50-51.
9. Braden B, Bergstrom N. Predictive validity of the Braden Scale for pressure sore risk in a nursing home population. *Research in Nursing & Health*. 1994;17:459-470.
10. Chang AM, Chau JPC, Holroyd E. Translation of questionnaires and issues of equivalence. *Journal of Advanced Nursing*. 1999;29(2):316-322.
11. Coutu-Wakulczyk G, Chartier L. French validation of the critical care family needs inventory. *Heart & Lung*. 1990;19(2):192-196.
12. Delgreco L, Walop W, Eastridge L. Questionnaire development: 2. Validity and reliability. *CMAJ*. 1987;136:699-700.
13. —. Questionnaire development: 3. Translation. *CMAJ*. 1987;136:817-818.
14. Keast D, Parslow N, Houghton PE, Norton L, Fraser C. Best practice recommendations for the prevention and treatment of pressure ulcers: Update 2006. *Wound Care Canada*. 2006;4(1):31-43.
15. Langemo DK, Olson B, Hunter S, Hanson D, Burd C, Cathcart-Silberberg T. Incidence and prediction of pressure ulcers in five patient care settings. *Decubitus*. 1991;4(3):25-6, 28, 30, 32, 36.
16. Lyder Ch, Yu C, Stevenson D, Mangat R, Empleo-Frazier O, Emerling J, Mckay J. Validating the Braden Scale for the prediction of pressure ulcer risk in blacks and latino/hispanic elders: A pilot study. *Ostomy/Wound Management*. 1998;44(3A Suppl):425-505.
17. Nelson McDermott MA, Palchanes K. A literature review of the critical elements of translation theory. *IMAGE: Journal of Nursing Scholarship*. 1994;26(2):113-117.
18. Registered Nurses' Association of Ontario (RNAO). *Nursing Best Practice Guideline: Risk Assessment and Prevention of Pressure Ulcers*. Toronto: RNAO. 2005.
19. —. *Nursing Best Practice Guideline: Assessment and Management of Pain*. Toronto: RNAO. 2002.
20. Robichaud-Ekstrand S, Haccoun RR, Millette D. Une méthode pour faire valider la traduction d'un questionnaire. *Canadian Journal of Nursing Research*. 1994;26(3):77-87.
21. Salvadlena GD, Snyder MI, Brogdon KE. Clinical trial of the Braden Scale on an acute care medical unit. *Journal of ET Nursing*. 1992;19(5):160-165.
22. Tourtual DM, Riesenber LA, Korutz CJ, Semo AH, Asef A, Talati K, Gill RDF. Predictors of hospital acquired heel pressure ulcers. *Ostomy/Wound Management*. 1997;43(9):24-40.
23. Varicchio CG. *Instruments for Clinical Health-Care Research*, Third Edition. Sudbury: Jones and Bartlett Publishers. 2004.

Systemic sclerosis *continued from page 18*

3. Aryal B, Khunder S, Schab E. Meta-analysis of systemic sclerosis and exposure to solvents. *American Journal of Industrial Medicine*. 2001;40:271-274.
4. Bovenzi M, Barbone F, Pisa FE, Betta A, Romeo L, Tonello A, Biasi D, Caramaschi P. A case-control study of occupational exposures and systemic sclerosis. *Int Arch Occup Environ Health*. 2004;77:10-16.
5. Hafner I, Schneider AE, Burg BG, Cassina PC. Management of leg ulcers in patients with rheumatoid arthritis or systemic sclerosis: The importance of concomitant arterial and venous disease. *J Vasc Surg*. 2000;32:322-9.
6. Sibbald RG, Williamson D, Orsted HL, Campbell K, Keast D, Krasner D, Sibbald D. Preparing the wound bed: Debridement, bacterial balance, and moisture balance. *Ostomy/Wound Management*. 2000;46:14-35.
7. Sandqvist G, Akesson A, Eklund M. Evaluation of paraffin bath treatment in patients with systemic sclerosis. *Disability and Rehabilitation*. 2004;26(16):981-987.
8. Pizzo G, Scardina GA, Messina P. Effects of a non-surgical exercise program on the decreased mouth opening in patients with systemic sclerosis. *Clin Oral Investig*. 2003;7:175-8.
9. Pfizenmaier D, Kavros S, Liedl D, Cooper L. Use of intermittent pneumatic compression for treatment of upper extremity vascular ulcers. *Angiology*. 2005;56:417-423.



It's not just silver. It's ACTICOAT silver.

Protect Yourself. Protect Your Patients
against a broad range of organisms
including MRSA to enhance wound healing.¹

The speed. The power. The trust.

1. Strohal R, et al. J Hosp Infect 2005;60:226-230. * Trademark of Smith & Nephew © 2006.
™SILCRYST is a trademark of NUCRYST Pharmaceutical Corp., used under license by Smith & Nephew Inc.

 **smith&nephew**
ACTICOAT[®]
Wound Dressings

AN ESSENTIAL COMPONENT OF
WOUND BED PREPARATION



150
years
Innovation.
For generations.