

# Puzzling Cases:

## Non-healing Venous Leg Ulcer



BY Rob Miller

**Rob Miller, MD, FRCPC,** has been practising dermatology for the past 20 years. He worked as a general practitioner in Ontario, British Columbia and South America before pursuing his studies in dermatology at McGill University in Montreal, QC. He is currently Associate Professor of Medicine at Dalhousie University and Co-director of the Chronic Wound Care Clinic at the QEII Hospital in Halifax, NS.

**A** 65-year-old female presents with a chronic venous ulcer (Figure 1) of one year's duration that will not heal with standard venous ulcer management, including adequate compression. Her Doppler studies are normal with respect to her arterial circulation, and her venous studies show deep-vein incompetence.



### Questions

1. Why does this ulcer not heal?
2. What clinical features suggest the cause of non-healing?
3. How would you restart the healing process?

### Answers

1. This ulcer illustrates the features of critical colonization. As shown in Diagram 1, there is a range of bacterial insult in a wound depending on not only the virulence of the organism but also the quantity of bacteria and the host resistance to this bacterial growth. A wound can normally tolerate contamination or colonization to a certain degree, but once critical colonization develops there is less chance for wound healing to occur.
2. Critical colonization can stall or create a non-healing


wound environment. Signs of critical colonization are as follows:

- (a) **N**on-healing wound
- (b) Increased **E**xudate
- (c) Increased **R**ed colour or easy friability or bleeding
- (d) Increased **D**ebris
- (e) Increased **S**mill

### Remember N.E.R.D.S.

3. The fastest way to kick-start the healing process is through debridement by means of curettage. Subsequent treatment of the wound with a silver dressing or cadexomer iodine would help to prevent critical recolonization. Curettage may have to be repeated more than once. ☹

DIAGRAM 1





## Now you can manage the wound...and the odor.

Most occlusive dressings do a good job of managing moisture, but unfortunately they can also contribute to wound odor. Now there's a new option...

### **Introducing Exuderm® OdorShield™—the first odor-control hydropolymer**

This patented new dressing uses cyclodextrins—naturally occurring, noncytotoxic molecules—to absorb odor as it absorbs wound exudate. It is the only dressing that combines the benefits of an advanced moisture management with powerful odor control.

### **More effective than charcoal**

The Exuderm OdorShield dressing outperformed leading charcoal-based dressings in laboratory odor absorption tests.\* While charcoal's odor-absorbing properties are deactivated by the proteins in wound exudate, Exuderm OdorShield's are actually enhanced. Other advantages: Exuderm OdorShield is conformable, absorbs more fluid than charcoal dressings, and requires no secondary dressing.

### **An improvement on yesterday's hydrocolloids**

Exuderm OdorShield eliminates the major drawback of traditional hydrocolloids: their distinctive odor upon removal. It effectively absorbs exudate and provides a protective, occlusive barrier that promotes wound healing. Features include a tapered edge and low-friction backing to prevent rollup and extend wear time.

**Ask your Medline representative for more information on Exuderm OdorShield dressings.**



**1-800-396-6996 | 905-403-7000**  
**www.medline.com**

\*Data on file.

©2006 Medline Industries, Inc.  
Exuderm and Medline are registered trademarks and OdorShield is a trademark of Medline Industries, Inc.