

A panel of 19 Physicians, NSWOCs, Wound Specialists, and Therapists with experience in treating VLUs, using the Muscle Pump Activator device, and advanced wound treatments.

This panel agreed that the geko™ device (Muscle Pump Activator) should be added to the treatment plan when:

- A patient cannot tolerate compression
- A patient is not in optimal compression
- No progress is seen in a wound after 2-4 weeks
- A wound has not healed 30% in 30 days

- Dr Asem Saleh
- Dr John Hwang
- Rosemary Hill
- Josee Senechal
- Michele Langille

- Bev Smith
- Carly St Michel
- Paulo da Rosa
- Amanda Loney
- Michele Labbie

- Sheri McPhee
- Allison Luther
- Maryse Beaumier
- Helen Arputhanathan
- Shannon Handfield

Read the VLU consensus here:  
<https://sites.google.com/view/VLUconsensus>  
or scan the QR code below



1. Bull R et al. Int Wound J. 2023; 1-9  
2. Jones N et al. Br J Nurs 2018; 27(20): S16-S21.

MDADWNCAN0772

[www.gekowound.ca](http://www.gekowound.ca)  
[info@perfusemedtec.com](mailto:info@perfusemedtec.com)

*The geko™ device demonstrated greater than two-fold increase in wound healing rate<sup>1</sup> and a reported reduction of pain<sup>2</sup> in venous leg ulcers vs compression alone.*  
**Harding et al, 2023**



Professor  
Keith Harding

Leads:



Dr Michael C. Stacey



Dr Robyn Evans



Dr Gary Sibbald

