

PluroGel®

Burn and Wound Dressing with concentrated surfactant (Micelle Matrix™) technology

- Poloxamer 188 activates wound healing
- Decreases inflammation & tissue damage¹
- Improves blood flow after burn injury²
- Blocks adhesion of certain proteins to prevent microbial adhesion³



IoPlex®

Iodophor Foam Dressing IoPlex® with I-Plexomer™ technology is the world's only controlled release iodine foam dressing

- Reduces bacterial burden within the wound dressing
- Effectively removes exudate and debris
- Released over 24 to 72 hours
- Can be cut to shape of wound and stacked
- Demonstrated a 4 log or greater kill against MRSA in 5 min and P. aeruginosa in 30 min.⁴



Medline develops innovative skin health products that can improve care outcomes.

¹Curry DJ, Wright DA, Lee RE, Kang UJ, Frim DM. Surfactant poloxamer 188-related decreases in inflammation and tissue damage after experimental brain injury in rats. *Journal Neurosurg* 2004;101(1 Suppl). Accessed November 29, 2018.

²Hunter RL, Luo AZ, Zhang R, Kozar RA, Moore FA. Poloxamer 188 inhibition of ischemia/reperfusion injury: evidence for a novel antiadhesive mechanism. *Ann Clin Lab Sci*. 2010;40(2):115-125.

³Birchenough SA, Rodeheaver GT, Morgan RF, Peirce SM, Katz AJ. Topical poloxamer-188 improves blood flow following thermal injury in rat mesenteric microvasculature. *Annals of Plastic Surgery*. Accessed November 29, 2018.

⁴Tharmalingam T, Ghebeh H, Wuerz T, Butler M. Pluronic enhances the robustness and reduces the cell attachment of mammalian cells. *Mol Biotechnol* 2008; 39(2):167-177.

⁵Data on file.

Together, we can transform injured skin into healthy skin

Discover transformative solutions designed to stimulate tissue regeneration by visiting Medline.ca or contacting your Medline Representative today at **1-800-268-2848**



**Skin
Health**